

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Ultrasound on the Frontlines of COVID-19: Report From an International Webinar

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1743442> since 2020-07-08T20:59:00Z

*Published version:*

DOI:10.1111/acem.14004

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

ACR guidelines on radiography in COVID-19 suggest portable CXR when it is considered “medically necessary”.<sup>3</sup> Further data is needed on the sensitivity, specificity, and diagnostic impact of CXR on suspected and diagnosed COVID-19 patients.

## Ultrasonography

Ultrasound of the lung utilizes artifacts and findings at the lung periphery, and early reports show that abnormal findings are common in COVID-19 patients. A series of twenty patients with COVID-19 described thickening and irregularity of the pleural line, a variety of B-line patterns and subpleural consolidations. Pleural effusions were rare.<sup>6</sup> A recent comparison of imaging modalities in intensive care patients suffering from acute respiratory failure demonstrated high agreement between lung ultrasound and CT. Ultrasound out-performed chest radiography particularly when assessing interstitial pathologies, as well as ground glass opacities and consolidations.<sup>6</sup> Since the sensitivity of CT is high in COVID-19 patients and progression of disease is apparent, it is likely that lung ultrasound closely mirrors the longitudinal changes found through CT. High sensitivity and specificity of lung ultrasound seen in acute respiratory distress syndrome (ARDS) as well as in H1N1 influenza suggest that similar test characteristics exist for lung ultrasound in COVID-19. Lung ultrasound also shows prognostic capabilities in ARDS before hypoxemia becomes evident.<sup>7</sup> While reports on lung ultrasound in COVID-19 are preliminary, they suggest ultrasound findings are likely more common than findings on plain chest radiography. POCUS may also identify and exclude other pulmonary causes of dyspnea as well as cardiac abnormalities. Future research should further specify test characteristics of ultrasound in COVID-19, as well as assessments of harm to ultrasound operators who risk increased contact with infected patients.

## ***Expert Observations from the Town Hall:***

### 1. Ultrasound in the diagnosis and monitoring of COVID-19

In the initial evaluation of a patient with suspected COVID-19, panelists posited that either a completely normal or a completely abnormal lung ultrasound may be helpful. A completely normal examination likely excludes a patient requiring further investigation at that time. Characteristically abnormal findings in a person under investigation for COVID-19 may identify patients requiring further evaluation or closer observation before RT-PCR results return. It is unclear whether B-line or consolidation thresholds exist that predict significant clinical deterioration in well-appearing patients who display lung findings. Observations that ultrasound findings may precede clinical symptoms suggests that ultrasound may identify more severe illness prior to the development of severe symptoms.

While lung ultrasound may be of little utility in patients who are critically ill from COVID-19, it may help exclude other pulmonary diseases (including pleural effusion or pneumothorax).<sup>6</sup> Incorporation of cardiac ultrasound could also identify cardiac complications from COVID-19 and it is recommended that a focused cardiac ultrasound be incorporated into the evaluation of symptomatic COVID-19 patients. Early reports suggest there may be direct cardiac ramifications of COVID-19 including gross LV dysfunction, elevated troponin without ST segment elevation, and potential RV

dilation.<sup>8,9</sup> Additional to lung ultrasound, cardiac views can help identify or exclude cardiac dysfunction and estimate fluid status through evaluation of the inferior vena cava.

Critically ill COVID-19 patients may present particular issues with vascular access that ultrasound assists with. Due to respiratory distress, some patients are unable to lie flat, making central venous access difficult. Ultrasound-guided vascular access was found to be quite helpful to establish multiple points of large bore peripheral access, in conjunction with intraosseous lines. Table 1 summarizes the main ultrasound applications used and their associated findings in COVID-19 patients.

Future research is needed to clarify the above associations between ultrasound findings and presence of illness, prognostication of illness severity, and ultimate patient outcomes. Panelists emphasized that while ultrasound findings are often characteristic of COVID-19 and may be more prominent in more severe disease, disposition decisions such as admission and level of care would likely be made predominantly on clinical appearance.

2. Ultrasound equipment

Newer pocket-carried ultrasound devices using tablet and smartphone technology may be easier to maneuver, protect, and clean after use on COVID-19 patients. It is feasible for the screen to be completely encased in a covering such as a Ziploc bag, and for transducers to be fully enclosed using sheaths. They may be particularly useful in situations where separate care or triage areas for cohorted COVID-19 patients are established outside the main areas where traditional machines reside. Additionally, teleguidance software built into these devices may be useful in providing remote instruction or evaluation of patients who are physically located away from typical treatment areas. Telemedicine and cloud-based image sharing may improve the capacity for care outside the hospital setting, and serial lung exams may provide objective data in remote or virtual clinical decision-making.

3. Infection control issues

Aggressive infection control was emphasized by all panelists, as a recent report suggested that the SARS-CoV-2 virus could survive for days on some surfaces.<sup>10</sup> Routine cleaning and disinfection procedures recommended by the Centers for Disease Control are particularly important when procedures with high risk of aerosolization are performed. Prior to entering the room, all unnecessary equipment including extra probes should be removed from the machine to minimize surface exposures. If available, transparent covers for machines and probes can be considered. If resources allow, equipment that is dedicated for use in patients with suspected or confirmed COVID-19 may be helpful. The Environmental Protection Agency maintains a list of SARS-CoV-2 approved disinfectants (<https://www.epa.gov/pesticide-registration/list-n-disinfectants-use-against-sars-cov-2>).

**Conclusion**

Ultrasound as a diagnostic test presents distinct advantages for imaging in COVID-19. It is a mobile technology that can be used in diverse environments, including in triage tents or makeshift hospitals that are now established for COVID-19 evaluation at many centers. POCUS in particular, where the clinician both performs and interprets the images immediately at the bedside, can minimize involvement of additional personnel in an infectious situation and provide immediate diagnostic information. However, care must be taken to avoid increased transmission of disease.

Incorporating ultrasound into the evaluation of COVID-19 patients will depend upon available resources, expertise of personnel, and logistic configurations unique to each situation. Further research and data are needed to determine the role of ultrasound as a screening tool for establishing both admission thresholds and level of care, its use in prognostication and monitoring of inpatients, as well as novel uses like home telemonitoring in discharged patients. Ultrasound appears promising as a first-line and perhaps comprehensive diagnostic imaging modality in suspected or diagnosed COVID-19.

The authors wish to acknowledge ACEP in arranging this webinar, as well as the Society of American Emergency Medicine, the American Academy of Emergency Medicine, and the Society of Clinical Ultrasound Fellowships for co-hosting this panel.

## References

1. Ai T, Yang Z, Hou H, Zhan C, Chen C, Lv W, *et al.* Correlation of Chest CT and RT-PCR Testing in Coronavirus Disease 2019 (COVID-19) in China: A Report of 1014 Cases [published online ahead of print February 26 2020]. *Radiology*. doi:10.1148/radiol.2020200642.
2. Pan F, Ye T, Sun P, Gui S, Liang B, Li L, *et al.* Time Course of Lung Changes On Chest CT During Recovery From 2019 Novel Coronavirus (COVID-19) Pneumonia [published online ahead of print February 13 2020]. *Radiology*. doi:10.1148/radiol.2020200370.
3. ACR Recommendations for the use of Chest Radiography and Computed Tomography (CT ) for Suspected COVID-19 Infection. (2020, March 22). Retrieved from <https://www.acr.org/Advocacy-and-Economics/ACR-Position-Statements/Recommendations-for-Chest-Radiography-and-CT-for-Suspected-COVID19-Infection>.
4. Ng MY, Lee E, Yang J, Yang F, Li X, Wang H, *et al.* Imaging Profile of the COVID-19 Infection: Radiologic Findings and Literature Review [published online ahead of print February 13 2020]. *Radiol Cardiothorac Imaging*. doi:10.1148/ryct.2020200034.
5. Pen QY, Wang XT, and Zhang LN. Findings of lung ultrasonography of novel corona virus pneumonia during the 2019 – 2020 epidemic [published online ahead of print March 12 2020]. *Intensive Care Med*. doi:10.1007/s00134-020-05996-6.
6. Tierney DM, Huelster JS, Overgaard JD, Plunkett MB, Boland LL, St. Hill CA, *et al.* Comparative performance of pulmonary ultrasound, chest radiograph, and CT among patients with acute respiratory failure. *Crit Care Med*. 2020;48(2):151-7.
7. Soldati G, Smargiassi A, Inchingolo R, Buonsenso D, Perrone T, *et al.* Is there a role for lung ultrasound during the COVID19 pandemic? [published online ahead of print March 20 2020]. *J Ultrasound Med*. doi:10.1002/jum.15284.

8. Arentz M, Yim E, Klaff L, Lokhandwala S, Riedo FX, Chong M, et al. Characteristics and Outcomes of 21 Critically Ill Patients With COVID-19 in Washington State [published online ahead of print March 19 2020]. *JAMA*. doi: 10.1001/jama.2020.4326.

9. Danzi GB, Loffi M, Galeazzi G, Gherbesi E. Acute pulmonary embolism and COVID-19 pneumonia: a random association? [published online ahead of print March 30 2020]. *Eur Heart J*. doi:10.1093/eurheartj/ehaa254.

10. VanDoremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *NEJM*. 2020;382(16):1564-7.

For Review Only

*Table 1: Point-of-Care Ultrasound Applications and Findings in COVID-19 patients*

| Lung Ultrasound   |   |
|---|---|
| <ul style="list-style-type: none"> <li>○ Panelists used 6-12 zones, or less</li> </ul>  | <ul style="list-style-type: none"> <li>● Thickened, irregular pleural line</li> <li>● B-lines with prominence in different locations (“patchy” appearance). B-lines are ring down artifacts from interstitial fluid characteristic of pneumonitis.</li> <li>● Subpleural consolidations</li> <li>● Larger consolidations</li> </ul> |
| Cardiac Ultrasound  |   |
| <ul style="list-style-type: none"> <li>○ Parasternal long and short axis</li> <li>○ Apical 4-chamber view</li> <li>○ Subxiphoid view</li> <li>○ Subxiphoid long axis (Inferior Vena Cava) view</li> </ul> | <ul style="list-style-type: none"> <li>● Pericardial effusion</li> <li>● Depressed ejection fraction or gross LV dysfunction</li> <li>● RV enlargement &amp; evidence of strain</li> <li>● Hypo- or hypervolemia</li> </ul>   |
| Vascular  |   |
|   | <ul style="list-style-type: none"> <li>● Deep venous thrombosis</li> <li>● Guided peripheral or central access</li> <li>● Guided arterial lines</li> </ul>  |

*NB: Panelists noted that there may be significant difficulties scanning all lung or cardiac views due to patient positioning or comfort, and levels of distress.*